

HFI systematics. Lessons learned

Guillaume Patanchon for the Planck
Collaboration

Lessons from Planck

- ❑ The data analysis and cleaning was a long process and required many iterations
- ❑ At the end, we reached the detector fundamental limit for cosmological channels
- ❑ Some effects were not expected at the level we found them in flight data
 - > ADC non-linearities
 - > Long time constants
 - > Response to cosmic rays
 - > 1/f noise
 - > Band-pass mismatch
- ❑ Coupling between effects was problematic. Ex: 4K lines and ADC non-linearities
- ❑ but for future experiment targeting $\sigma_r < 10^{-3}$, systematic effects must be controlled to a higher precision, although many effects will probably scale as 1/Ndet.
- ❑ Importance of observation redundancies: different survey, different scanning angle (limited for Planck), different detectors etc..., **importance of the dipole**, 353 GHz is harder to process
- ❑ Importance of house keeping data. E.g: fully sampled raw data for the ADC correction.
- ❑ Many affect as band-pass mismatch, polarization efficiency, calibration are coupled and need to be corrected at the map-making level, with the help of the dipole

Data reduction

Model of the raw data:

$$d_i(t) = \underset{\text{Gain}}{g_i} \int R_i(t - t') \underset{\text{Transfer function}}{W(t')} \left[X_i(t') + \sum_j \underset{\text{4K lines } (A_k, w_k, \dots)}{T_{ij}(t')} \right] dt' + Q_i(t) + n_{J_i}(t) + \sum_c F_{ic}(t).$$

$$X_i(t') = \left[\int \underset{\text{Lobes}}{H_i(t' - t'')} \left(\{B_{i;\psi_{it''}} * [S_i + o]\}(r_{t''}^{\vec{r}}) + n_{s_i}(t'') \right) dt'' \right]$$

Data are digitized, averaged over 40 samples, and compressed on board

Data processing: compression




Goal to reach

$$d_i(t_p) = \{B_{\psi_{it_p}} * [S_i + o]\}(r_{t_p}^{\vec{r}}) + n_{i;\text{total}}(t_p).$$



Symmetrized lobe

Cosmic rays at L2



Mainly galactic protons and Helium nuclei


CR of ~ 1 GeV dominate



Cut off due to material around the detectors at ~ 50 MeV

No contribution from solar particles which can not reach the detectors, except during flares


Amplitude of the spectrum at L2 is modulated by solar activity




Thermal fluctuations of the focal plate are caused by CR:

Compensated by PIDs on time scales > 1 h

CR interaction with HFI detectors



Thermal modeling is important. Long time constants come from the links between the wafer and the detector housing




- Long glitches are direct impact of protons in the silicon wafer

- short glitches are direct impact of protons in the grid/thermistor. Should be representative of response to photons.

This was proved with the help of ground tests with alpha particles

Ground tests and thermal modeling



Ground tests did not provide a definitive answer on the thermal path



Basic Equation

$$C_j \frac{dT_j}{dt} = \sum_{i=0}^{12} G_{ij} (T_i - T_j)$$



Simulation of a 23MeV Proton in the silicon die



Cosmic ray removal

Joint fit of templates for each detected event.


- Removal of long glitch tails
- Flagging 10 to 25 % of data depending on the detector




Analysis made difficult because of the high confusion of events

Residual at the level of noise for the worst channels at low frequencies < 0.2 Hz

At the end, the glitch contribution to the noise on the maps is significant only for $\ell < 10$, still smaller than detector noise



Noise in HFI time ordered data





Glitches below the detection threshold common between PSB-a and PSB-b
Provide a limit on the level of remaining glitches in data

Lesson learned

CR signals were a rich probe of detector and focal plane parameters, allowing to constrain some systematic effects

- Thermal links in the focal plane



- Coherent picture of the interaction: Ballistic phonons + thermal propagation


- Origin of the low frequency noise, correlated among detectors

➡ Events on the bolometer plate


- Long time constants

The 2-second TC induces a bias on the power spectrum of $\sim 1-2\%$ if uncorrected


Dipole shift depending on the scan direction



Lessons for future experiments



- All events were detected in Planck!!
- Ballistic phonons helped the detection
- Could create correlated noise with large wafers



Main systematic effects


- Additive effects : Glitches, unexpected 1/f noise, microphonic noise
- Main effects I to P leakages, different detectors had to be combined to estimate Q and U Stokes parameters
 - ADC non-linearities
 - Band-pass mismatch
 - Long time constants
- Other systematics
 - Beam + time constants
- Use of redundancies of observations and of the strong dipole signal to calibrate and correct the data : **Surveys with opposite scanning directions allowed optimization of parameters and correction of many systematic effects.**

Many effects scale with $\langle \cos 2\Psi \rangle$ and $\langle \sin 2\Psi \rangle$. The use of a HWP and better angle redundancies as planned for LiteBIRD help.

Survey difference maps

Survey difference maps were useful to track and characterize systematic effect

217GHz I map NO VLTC CORRECTED S1-S2



Uncorrected **long time constants**
slightly shift the galaxy
Residual dipole seen in the difference

-10.0  10.0 microK




Corrected after optimization at the
map-making level by template
fitting

Beam and transfer function estimation

- Time response is degenerate with the beam response
- The time response and beam shapes are estimated using a combination of planet scans (by symmetrizing the beam shape), galaxy crossings, bias steps (CPV phase) and glitch data.
- The pointing uncertainties (~ 3 arcsec) and glitch is the main source of errors in the main lobe estimation



Corrections of the transfer function at the likelihood optimization stage



ADC non-linearities

Analog to Digital convertor have some non-linearities





Cosmological data Planck data are sampling -300 to +300 ADUs near the code 32768, that's where the non-linearity is larger! + drifts of data

Uncertainties in the 4K line freqs, coupled with ADC



Main systematic effect in Planck for polar.


Electronic response



House-keeping data were essential!


ADC correction

The correction is very effective but limited by the 4K line estimation.



A second correction was performed at the map-making level

Jackknife : positive – negative parities




ADC is the limiting systematic effect in Planck for polarization measurement

No intrinsic detector gain variations have been detected

Band-pass mismatch


Differences in the band shapes from detector to detector induced intensity to polarization of galactic components when calibrating on CMB



CO transition line 1- \rightarrow 0 falls at the edge of the 100 Hz filters so the CO components has very different amplitude from detector to detector

After integrating the dust spectrum:

A few percent effects for the amplitude of the dust from detector to detector




Band-pass mismatch correction

-Band passes were measured from the ground, but leakage coefficients have to be estimated from flight data

$$m = T_{Sky} + (\gamma_{Dust} - 1)T_{Dust} + (\gamma_{CO} - 1)T_{CO} + \dots$$

- **Joint estimation of CO and dust leakages at the map-making level.** Naturally minimizes the survey difference contamination. Coupled with many effects.



Effect mostly removed at the end

Summary of systematic effects (HFI)

- ADC is the dominant systematic effect
- Its contribution is at the level of the noise at low l s
- Other systematic effects are negligible after processing

